

Analysis of the $\text{Sn}^{7+}\dots^{14+}$ fine structure by optical and ultraviolet spectroscopy in an EBIT

F. Torretti^{1,2}, A. Windberger^{1,3}, A. Borschevsky⁴, A. Ryabtsev⁵, S. Dobrodey³, R. Hoekstra^{1,6}, W. Ubachs^{1,3},

J. R. Crespo López-Urrutia³, and O.O. Versolato¹

¹ EUV Plasma Dynamics, Advanced Research Center for Nanolithography (ARCNL), Science Park 110, 1098 XG Amsterdam, The Netherlands

² Department of Physics and Astronomy, LaserLaB, Vrije Universiteit, De Boelelaan 1081, 1081 HV Amsterdam, The Netherlands

³ Max-Planck-Institute for Nuclear Physics, Saupfercheckweg 1, 69117 Heidelberg, Germany


⁴ Van Swinderen Institute, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

⁵ Institute of Spectroscopy, Russian Academy of Sciences, Troitsk, Moscow oblast, 142190 Russia


⁶ Zernike Institute for Advanced Materials, University of Groningen, Nijenborgh 4, 9747 AG Groningen, The Netherlands

Highly charged ions of tin (Sn^{7+} - Sn^{14+}) are of technological interest as these are used in laser-produced-plasma (LPP) sources for the generation of extreme ultraviolet (EUV) light for nanolithographic applications. Here, we present the results from spectroscopic measurements of these ions, both in the EUV and optical regimes, obtained in a charge-state-resolved manner with the FLASH-EBIT (Electron Beam Ion Trap) at the Max Planck Institute for Nuclear Physics in Heidelberg. Transitions were identified by the use of *ab initio* Fock space coupled cluster calculations, thus confirming the predictive power of this tool. Our identifications were further corroborated using semi-empirical calculations within the Cowan code framework. With our experimental and theoretical work we have re-evaluated the fine structure of Sn^{11+} - $^{14+}$, and comparison with previous, foundation-laying, studies suggest that some of their level identifications of the Sn HCl atomic structure based on EUV spectral data need to be revisited [1].

Charge-state-resolved spectra



The atomic structure of Sn ions



[Kr] $4d^{m-1}4f$
[Kr] $4d^{m-1}5p$
[Ar] $3d^{10}4s^24p^54d^m$
[Ar] $3d^{10}4s^24p^54d^m5s$


13.5 nm

[Ar] $3d^{10}4s^24p^64d^m$


State density diagram
 $\text{Sn}^{7+}\dots^{14+}$ ions contribute strong electric dipole transitions to an unresolved transition array (UTA). In the optical regime M1 transitions can be observed, directly connecting the ground state structure of these ions. In this regime, only few tens of transitions are present which can be resolved with ease, compared to the EUV regime.

Charge state identification (Sn^{7+} - Sn^{10+})

The EBIT EUV spectra are compared with known lines [2] to identify the emissions by the lower charge states of Sn in the optical regime.




Charge state identification (Sn^{11+} - Sn^{14+})




Fluorescence curves are used to assign lines to their respective charge states. The double-peaked structure of these curves and the fact that the fluorescence starts before the ionization potentials are strong evidence of metastable states populations inside the EBIT.

Fluorescence curves and metastable states



Comparison with theoretical calculations

Identifying optical transitions with the help of theory: Fock space coupled cluster [4,5] and semi-empirical calculations within the COWAN code framework [6]. This data enables a reinterpretation of the ground state configurations in the studied ions.



Conclusions

- 120 M1 transitions have been identified in the Sn ions of relevance for EUV sources;
- Our experimental input indicates the need for re-interpretation of foundation-laying work on the atomic structure of Sn ions (in Sn^{11+} - Sn^{14+});
- The majority of energy levels in the ground configurations of Sn^{8+} and Sn^{9+} have been determined;
- The predictive power of FSCC has been shown by the agreement with the experimental data.

References

- [1] A. Windberger et al, Phys. Rev. A 94, 012506 (2016).
- [2] S. S. Churilov and A. N. Ryabtsev, Physica Scripta 73, 614 (2006).
- [3] S. S. Churilov and A. N. Ryabtsev, Optics and Spectroscopy 101, 169 (2006).
- [4] D. Mukherjee and S. Pal, Adv. Quantum Chem. 20, 291 (1989).
- [5] U. Kaldor and E. Eliav, Adv. Quantum Chem. 31, 313 (1998).
- [6] A. N. Ryabtsev, E. Y. Kononov, and S. S. Churilov, Optics and Spectroscopy 105, 844 (2008).



ADVANCED RESEARCH CENTER FOR NANOLITHOGRAPHY



MAX-PLANCK-INSTITUT FÜR KERNPHYSIK

ARCNL is a public-private partnership between FOM, UvA, VU and ASML

